Contribution ID: 80 Type: Rapid

## High Temperature Proximity Effect in Topological Insulators

Thursday, 26 July 2012 16:45 (12 minutes)

Kenneth Burch1, Parisa Zareapour1, Alex Hayat1,2, Shu Yang F. Zhao1, Anjan Reijnders1, Mikhail Kreshchuk1, Achint Jain1, Daniel C. Kwok3, Nara Lee3, Sang-Wook Cheong3, Zhijun Xu4, Alina Yang4, G. D. Gu4, and R. Cava5

 $1 Department of Physics and Institute for Optical Sciences, University of Toronto, 60 \, St. \,\, George \,\, Street, Toronto \,\, ON, \,M5S \,\, 1A7, \, Canada$ 

2Centre for Quantum Information and Quantum Control University of Toronto, 60 St. George Street, Toronto ON, M5S 1A7, Canada

3Rutgers Center for Emergent Materials and Department of Physics and Astronomy, Rutgers University, 136 Frelinghuysen Road, Piscataway, New Jersey 08854, USA 4CMP&MS Department, Brookhaven National Laboratory, Upton, New York 11973, USA

5 Department of Chemistry, Princeton University, New Jersey 08544, USA

Producing new effects through the combination of different materials has a long history in science and technology. One of the most intriguing recent ideas is the emergence of Majorana fermions when a topological insulator is placed in proximity with a superconductor. Towards this goal, we produced high-temperature superconductivity in topological insulators via the proximity to a high temperature superconductor. In this talk I will describe the simple mechanism to achieve this. In addition I will present our extensive tunnelling spectroscopy studies that confirm the existence of the proximity effect. Lastly I will discuss some potential future directions this technique offers for novel optical devices.

**Primary author:** BURCH, Kenneth (University of Toronto)

Presenter: BURCH, Kenneth (University of Toronto)

**Session Classification:** High Tc Cuprates & Pnictides

Track Classification: Topological Insulators