CIPANP 2018 - Thirteenth Conference on the Intersections of Particle and Nuclear Physics

Contribution ID: 96 Type: Parallel

Search for $K^+ \to \pi^+ \nu \nu$ at CERN

Tuesday, 29 May 2018 15:20 (20 minutes)

The decay $K^+ \to \pi^+ \nu \nu$, with a very precisely predicted branching ratio of less than 10^{-10} , is one of the best candidates to reveal indirect effects of new physics at the highest mass scales. The NA62 experiment at the CERN SPS is designed to measure the branching ratio of the $K^+ \to \pi^+ \nu \nu$ with a decay-in-flight technique, novel for this channel. NA62 took data in 2016, 2017 and another year run is scheduled in 2018. Statistics collected in 2016 allows NA62 to reach the Standard Model sensitivity for $K^+ \to \pi^+ \nu \nu$, entering the domain of 10^{-10} single event sensitivity and showing the proof of principle of the experiment. The analysis data is reviewed and the preliminary result from the 2016 data set presented.

E-mail

bob.velghe@triumf.ca

Collaboration name

NA62

Primary author: Dr VELGHE, Bob (TRIUMF)

Presenter: Dr VELGHE, Bob (TRIUMF)

Session Classification: HFCKM / PPHI

Track Classification: PPHI