

Diagnosing New Physics with LUV and LFV B Decays

Alakabha Datta

University of Mississippi

May 29, 2018

CIPANP


Outline of Talk

- In recent times there have been some anomalies in B decays that indicate lepton non-universal new physics.
- These are in semileptonic $b \rightarrow c\tau\bar{\nu}_\tau$ transitions: $R_{D^{(*)}}$ puzzle.
- These are in semileptonic $b \rightarrow s\ell^+\ell^-$ ($\ell = \mu, e$) transitions: $R_K, R_{K^{(*)}}$ puzzles. BR of $b \rightarrow s\mu^+\mu^-$ modes are lower and also deviation in P'_5 angular observable.
- These all indicate LUV New Physics.

Plan of the Talk

- If NP is present how to probe this NP in distributions and related decays.
- LUV can often lead to lepton flavor violation.
- Will consider simultaneous explanation of $R_{D^{(*)}}$ and R_K puzzles ([1412.7164](#), [1609.09078](#)) and LFV tests .


$R_{D^{(*)}}$ puzzle



$$A_{SM} = \frac{G_F}{\sqrt{2}} V_{cb} \left[\langle D^{(*)}(p') | \bar{c} \gamma^\mu (1 - \gamma_5) b | \bar{B}(p) \rangle \right] \bar{\tau} \gamma_\mu (1 - \gamma_5) \nu_\tau$$

$$R(D) \equiv \frac{\mathcal{B}(\bar{B} \rightarrow D^+ \tau^- \bar{\nu}_\tau)}{\mathcal{B}(\bar{B} \rightarrow D^+ \ell^- \bar{\nu}_\ell)} \quad R(D^*) \equiv \frac{\mathcal{B}(\bar{B} \rightarrow D^{*+} \tau^- \bar{\nu}_\tau)}{\mathcal{B}(\bar{B} \rightarrow D^{*+} \ell^- \bar{\nu}_\ell)}$$

$R_D, R_{D^*}, \text{HFAG}$



Experiments: $R_{D^{(*)}}$ puzzle

The average of $R(D)$ and $R(D^*)$ measurements evaluated by the Heavy-Flavor Averaging Group are

$$R(D)_{exp} = 0.407 \pm 0.039 \pm 0.024, \quad (1)$$

$$R(D^*)_{exp} = 0.304 \pm 0.013 \pm 0.007. \quad (2)$$

The combined analysis of $R(D)$ and $R(D^*)$, taking into account measurement correlations, finds that the deviation is at the level of 4.1σ from the SM prediction.

$$R(D)_{SM} = 0.298 \pm 0.003,$$

$$R(D^*)_{SM} = 0.255 \pm 0.004. \quad (3)$$

There are lattice QCD predictions for the ratio $R(D)_{SM}$ in the Standard Model that are in good agreement with one another,

$$R(D)_{SM} = 0.299 \pm 0.011 \quad [\text{FNAL/MILC}],$$

$$R(D)_{SM} = 0.300 \pm 0.008 \quad [\text{HPQCD}].$$

Model independent NP analysis (See for example: Datta, Duraisamy, Ghosh)

At the m_b scale: $SU(3)_c \times U(1)_{em}$.


- Effective Hamiltonian for $b \rightarrow c l^- \bar{\nu}_l$ with Non-SM couplings. The NP has to be LUV.

$$\mathcal{H}_{\text{eff}} = \frac{4G_F V_{cb}}{\sqrt{2}} \left[(1 + V_L) [\bar{c} \gamma_\mu P_L b] [\bar{l} \gamma^\mu P_L \nu_l] + V_R [\bar{c} \gamma^\mu P_R b] [\bar{l} \gamma_\mu P_L \nu_l] \right. \\ \left. + S_L [\bar{c} P_L b] [\bar{l} P_L \nu_l] + S_R [\bar{c} P_R b] [\bar{l} P_L \nu_l] + T_L [\bar{c} \sigma^{\mu\nu} P_L b] [\bar{l} \sigma_{\mu\nu} P_L \nu_l] \right]$$

The NP can be probed via distributions and other related decays.

$B \rightarrow D^{(*)} \tau \nu_\tau$ in SM

The helicity amplitudes and consequently the NP couplings can be extracted from an angular distribution and compared with models.



Distributions have been measured very well by Belle for $B \rightarrow D^{(*)} l \nu_l$. We can then extract the Form Factors assuming no NP in these modes.

If we observe τ decay then we can measure τ polarization and CPV.

$B \rightarrow D^{(*)} \tau \nu_\tau$ in SM + NP, Helicity Amplitudes

Decay Distribution described by Helicity Amplitudes

$$\mathcal{H}_0 = \frac{4G_F V_{cb}}{\sqrt{2}} \frac{1}{2m_{D^*} \sqrt{q^2}} \left[(m_B^2 - m_{D^*}^2 - q^2)(m_B + m_{D^*}) A_1(q^2) - \frac{4m_B^2 |p_{D^*}|^2}{m_B + m_{D^*}} A_2(q^2) \right] (1 + V_L - V_R),$$

$$\mathcal{H}_{\parallel} = \frac{4G_F V_{cb}}{\sqrt{2}} \sqrt{2} (m_B + m_{D^*}) A_1(q^2) (1 + V_L - V_R),$$

$$\mathcal{H}_{\perp} = -\frac{4G_F V_{cb}}{\sqrt{2}} \sqrt{2} \frac{2m_B V(q^2)}{(m_B + m_{D^*})} |p_{D^*}| (1 + V_L + V_R),$$

$$\mathcal{H}_t = \frac{4G_F V_{cb}}{\sqrt{2}} \frac{2m_B |p_{D^*}| A_0(q^2)}{\sqrt{q^2}} (1 + V_L - V_R),$$

$$\mathcal{H}_P = -\frac{4G_F V_{cb}}{\sqrt{2}} \frac{2m_B |p_{D^*}| A_0(q^2)}{(m_b(\mu) + m_c(\mu))} (S_R - S_L).$$

Distributions

- $F_L (D^*)$ polarization. Distribution in θ^* .
- A_{FB} for both D and D^* . Distribution in θ_l .
- If we make the τ decay then we can measure the longitudinal tau polarization $P_\tau(D^{(*)})$.
- Finally we can look at CP violating terms in the angular distribution.

CPV Triple products






- There are triple products that appear in the angular distributions proportional to $\sin \chi$ (Datta and Duraisamy.)
- The triple product in the B rest frame: $\sim (\vec{n}_D \times \vec{n}_l) \cdot \vec{p}_{D^*} \sim \sin \chi$ with $\vec{n}_D \sim \vec{p}_D \times \vec{p}_\pi$ and $\vec{n}_l \sim \vec{p}_l \times \vec{p}_\nu$.
- These T.P. are proportional to $\mathcal{I}(H_i H_\perp^*)$. There are CPV. In the SM these terms are absent because all SM amplitudes have the same weak phase - V_{cb} .
- Since the p_τ momentum is not known we make the τ decay: $\tau \rightarrow V \nu_\tau$ and use the V momentum to construct the T.P. (Hagiwara, Nojiri, Sakaki).

Other Decays

NP can be constrained from other decays have the same quark transition as $R_{D^{(*)}}$

- $B_c \rightarrow \tau^- \bar{\nu}_\tau$ (Alonso, Grinstein, Camalich). $\Gamma[B_c] > \Gamma[B_c \rightarrow \tau^- \bar{\nu}_\tau]$. g_P coupling is very constrained.
- $B_c \rightarrow J/\psi \tau^- \bar{\nu}_\tau$ LHCb measurement finds about a 2σ deviation from the SM.
- $b \rightarrow \tau \nu X$ (LEP) (Saeed Kamali, AD).
- Measurements in $\Lambda_b \rightarrow \Lambda_c \tau \bar{\nu}_\tau$ can further constrain the NP parameter space. (Datta:2017aue, Shivashankara:2015cta).
- $\Lambda_b \rightarrow \Lambda_c$ form factors are calculated from lattice QCD (Datta:2017aue, Detmold:2015aaa)

$$R_{\Lambda_c}^{Ratio} = 1.3 \pm 3 \times 0.05$$



Interesting Facts



$$R(D)^{Ratio} = \frac{R(D)_{exp}}{R(D)_{SM}} = 1.36 \pm 0.15(1.30 \pm 0.17),$$
$$R(D^*)^{Ratio} = \frac{R(D^*)_{exp}}{R(D^*)_{SM}} = 1.19 \pm 0.06(1.25 \pm 0.08). \quad (7)$$

- If NP is just $V - A$ then


$$R_D^{ratio} \equiv \frac{R_D^{expt}}{R_D^{SM}} = |1 + V_L|^2 = R_{D^*}^{ratio} \equiv \frac{R_{D^*}^{expt}}{R_{D^*}^{SM}}.$$

- If NP couples to RH particles only

$$R_D^{ratio} \equiv \frac{R_D^{expt}}{R_D^{SM}} = (1 + |V_L|^2) = R_{D^*}^{ratio} \equiv \frac{R_{D^*}^{expt}}{R_{D^*}^{SM}}.$$

W' models from $SU(2)_L \times SU(2)_V \times U(1)_X \rightarrow SU(2)_L \times U(1)_Y$ (
1804.04135, 1804.04642

$b \rightarrow s\mu^+\mu^-$ Anomaly



$$H_{\text{eff}}(b \rightarrow s\ell\bar{\ell}) = -\frac{\alpha G_F}{\sqrt{2}\pi} V_{tb} V_{ts}^* [C_9 (\bar{s}_L \gamma^\mu b_L) (\bar{\ell} \gamma_\mu \ell) + C_{10} (\bar{s}_L \gamma^\mu b_L) (\bar{\ell} \gamma_\mu \gamma^5 \ell)] ,$$

$$H_{\text{eff}}(b \rightarrow s\nu\bar{\nu}) = -\frac{\alpha G_F}{\sqrt{2}\pi} V_{tb} V_{ts}^* C_L (\bar{s}_L \gamma^\mu b_L) (\bar{\nu} \gamma_\mu (1 - \gamma^5) \nu) ,$$

$$H_{\text{eff}}(b \rightarrow s\gamma^*) = C_7 \frac{e}{16\pi^2} [\bar{s} \sigma_{\mu\nu} (m_s P_L + m_b P_R) b] F^{\mu\nu}$$

R_K puzzle, Ratios of $b \rightarrow s\mu^+\mu^-$ and $b \rightarrow se^+e^-$. Part II (Clean), 1708.02515

$$R_K \equiv \mathcal{B}(B^+ \rightarrow K^+\mu^+\mu^-)/\mathcal{B}(B^+ \rightarrow K^+e^+e^-)$$

$$R_K^{\text{expt}} = 0.745^{+0.090}_{-0.074} (\text{stat}) \pm 0.036 (\text{syst})$$

$$1 \leq q^2 \leq 6.0 \text{ GeV}^2$$




Figure: Comparison of the measurements of R_K from LHCb (black dots), BaBar (red squares) and Belle (blue triangles) with the SM expectation (purple line).





Figure: Comparison of the measurements of R_{K^*} from LHCb with (left) SM predictions and (right) BaBar and Belle.

$$R_{K^*}^{\text{expt}} = \begin{cases} 0.660_{-0.070}^{+0.110} \text{ (stat)} \pm 0.024 \text{ (syst)} & 0.045 \leq q^2 \leq 1.1 \text{ GeV}^2, \\ 0.685_{-0.069}^{+0.113} \text{ (stat)} \pm 0.047 \text{ (syst)} & 1.1 \leq q^2 \leq 6.0 \text{ GeV}^2. \end{cases}$$

R_K and R_{K^*} in the SM very close to 1 in the central bin and
 $R_{K^*} \sim 0.92$ in the low bin.

$R_{K^{(*)}}$ puzzle: Other Experiment

- Measurements from Belle finds difference in same q^2 bin as LHCb

$$Q_5 = P'_5(\mu\mu) - P'_5(ee)$$


(1612.05014). Large errors.

- Low q^2 dominated by photon pole which is not LUV. Hence measurement difficult to understand with heavy NP.

Deviations in $b \rightarrow s\mu^+\mu^-$ Part I- Hadronic Uncertainty

- Anomalies appear in $B \rightarrow K^{(*)}\mu^+\mu^-$ (LHCb, Belle, Atlas, CMS) : Deviations branching ratios and in the angular observable like P'_5 .
- BR are lower than the SM predictions.
- (LHCb) $B_s^0 \rightarrow \phi\mu^+\mu^-$ which are lower than SM predictions based on lattice QCD and QCD sum rules.
- Note all these are in $b \rightarrow s\mu^+\mu^-$ and the SM predictions are not free of hadronic uncertainties.

P'_5 in $B \rightarrow K^*(K\pi)\mu^+\mu^-$






P'_5 in $B_d^0 \rightarrow K^* \mu^+ \mu^-$

$$\begin{aligned}
 & \frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d^4(\Gamma + \bar{\Gamma})}{dq^2 d\vec{\Omega}} \\
 &= \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_k + F_L \cos^2 \theta_k \right. \\
 &\quad + \frac{1}{4}(1 - F_L) \sin^2 \theta_k \cos 2\theta_l \\
 &\quad - F_L \cos^2 \theta_k \cos 2\theta_l + S_3 \sin^2 \theta_k \sin^2 \theta_l \cos 2\phi \\
 &\quad + S_4 \sin 2\theta_k \sin 2\theta_l \cos \phi + S_5 \sin 2\theta_k \sin \theta_l \cos \phi \\
 &\quad + \frac{4}{3} A_{\text{FB}} \sin^2 \theta_k \cos \theta_l + S_7 \sin 2\theta_k \sin \theta_l \sin \phi \\
 &\quad \left. + S_8 \sin 2\theta_k \sin 2\theta_l \sin \phi + S_9 \sin^2 \theta_k \sin^2 \theta_l \sin 2\phi \right]. \tag{8}
 \end{aligned}$$

Optimal observables. When E_K is large, small q^2 , in leading order in SCET these observables are free from form factors. Corrections are $\sim O(\frac{1}{E_K})$ and α_s .

$$\begin{aligned}
 P_1 &= \frac{2 S_3}{(1 - F_L)} = A_T^{(2)}, \\
 P_2 &= \frac{2 A_{FB}}{3(1 - F_L)}, \\
 P_3 &= \frac{-S_9}{(1 - F_L)}, \\
 P'_{4,5,8} &= \frac{S_{4,5,8}}{\sqrt{F_L(1 - F_L)}}, \\
 P'_6 &= \frac{S_7}{\sqrt{F_L(1 - F_L)}}.
 \end{aligned} \tag{9}$$

Just like $B \rightarrow D^{(*)} \tau \nu_\tau$ one can look at other observables like F_L, A_{FB} and CP violating co-efficients.



Recent Fits after $R_{K^{(*)}}$

Fits by many authors(1704.05435, 1704.05438, 1704.05444, 1705.05446, 1704.05447....) to all $b \rightarrow s\ell\ell$ observables: arXiv:1704.07397 : Alok et.al.

Scenario	WC	pull
(I) $\Delta C_9^{\mu\mu}(\text{NP})$	-1.25 ± 0.19	5.9
(II) $\Delta C_9^{\mu\mu}(\text{NP}) = -\Delta C_{10}^{\mu\mu}(\text{NP})$	-0.68 ± 0.12	5.9
(III) $\Delta C_9^{\mu\mu}(\text{NP}) = -\Delta C_9^{\prime\mu\mu}(\text{NP})$	-1.11 ± 0.17	5.6

Here NP effects only the muons.

Remember in the $R_{D^{(*)}}$ puzzle also indicated LH NP interactions. This gives a hint to connect the two anomalies.

LFV from LUV

- Glashow, Guadagnoli and Lane (GGL), 1411.0565 pointed out in general

LUV \Rightarrow LFV.

$$\frac{G}{\Lambda_{NP}^2} (\bar{b}'_L \gamma_\mu b'_L) (\bar{\tau}'_L \gamma^\mu \tau'_L) ,$$

where $G = O(1)$, $G/\Lambda_{NP}^2 \ll G_F$

- When one transforms to the mass basis, this generates the operator $(\bar{b}_L \gamma_\mu s_L) (\bar{\mu}_L \gamma^\mu \mu_L)$ that contributes to $\bar{b} \rightarrow \bar{s} \mu^+ \mu^-$.
The contribution to $\bar{b} \rightarrow \bar{s} e^+ e^-$ is much smaller, leading to a violation of lepton flavor universality.
- GGL's point was that LFV decays, such as $B \rightarrow K \mu e$, $K \mu \tau$ and $B_s^0 \rightarrow \mu e$, $\mu \tau$, are also generated.

R_K and $R_{D^{(*)}}$

Assuming the scale of NP is much larger than the weak scale, the semileptonic operators should be made invariant under the full $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge group. (Bhattacharya, Datta, London, Shivshankara, 1412.7164) considered two possibilities for LH interactions:

$$\begin{aligned}\mathcal{O}_1^{NP} &= \frac{G_1}{\Lambda_{NP}^2} (\bar{Q}'_L \gamma_\mu Q'_L) (\bar{L}'_L \gamma^\mu L'_L), \\ \mathcal{O}_2^{NP} &= \frac{G_2}{\Lambda_{NP}^2} (\bar{Q}'_L \gamma_\mu \sigma^I Q'_L) (\bar{L}'_L \gamma^\mu \sigma^I L'_L) \\ &= \frac{G_2}{\Lambda_{NP}^2} \left[2(\bar{Q}'_L{}^i \gamma_\mu Q'^j_L) (\bar{L}'_L{}^j \gamma^\mu L'^i_L) - (\bar{Q}'_L \gamma_\mu Q'_L) (\bar{L}'_L \gamma^\mu L'_L) \right].\end{aligned}$$

Here $Q' \equiv (t', b')^T$ and $L' \equiv (\nu'_\tau, \tau')^T$. The key point is that \mathcal{O}_2^{NP} contains both neutral-current (NC) and charged-current (CC) interactions. The NC and CC pieces can be used to respectively explain the R_K and $R_{D^{(*)}}$ puzzles.

UV completion

- UV completions considered by many authors e.g. [L. Calibbi, A. Crivellin and T. Ota, 1506.02661](#) considered possible UV completions that can give rise to $\mathcal{O}_{1,2}^{NP}$.
- (i) a vector boson (VB) that transforms as $(\mathbf{1}, \mathbf{3}, 0)$ under $SU(3)_C \times SU(2)_L \times U(1)_Y$, as in the SM.
- (ii) an $SU(2)_L$ -triplet scalar leptoquark (S_3) $[(\mathbf{3}, \mathbf{3}, -2/3)]$.
- (iii) an $SU(2)_L$ -singlet vector leptoquark (U_1) $[(\mathbf{3}, \mathbf{1}, 4/3)]$.
- $SU(2)_L$ -triplet vector leptoquark (U_3) $[(\mathbf{3}, \mathbf{3}, 4/3)]$.
- The vector boson generates only \mathcal{O}_2^{NP} , but the leptoquarks generate particular combinations of \mathcal{O}_1^{NP} and \mathcal{O}_2^{NP} .

Models

- Note to simply explain $b \rightarrow s\ell^+\ell^-$ we can have Z' $(\mathbf{1}, \mathbf{1}, 0)$ from $U(1)$. One can consider both $(\mathbf{1}, \mathbf{3}, 0)$ and $(\mathbf{1}, \mathbf{1}, 0)$.
- Models with $U(2)_q \times U_l(2)$ flavor symmetry and breaking: See for example: [Dario Buttazzo, Admir Greljo, Gino Isidori David Marzocca \(Zurich U.\) 1706.07808](#).
- Many of the general features can be understood in a simple analysis.
- In models other processes get affected and so specific models are more constrained.

Models: Bhattacharya, Datta, Guevin, London, Watanabe, 1609.09078

Models: **Vector Bosons and Leptoquarks.**

Transform to the mass basis:

$$u'_L = Uu_L, \quad d'_L = Dd_L, \quad \ell'_L = L\ell_L, \quad \nu'_L = L\nu_L,$$

The CKM matrix is given by $V_{CKM} = U^\dagger D$. The assumption is that the transformations D and L involve only the second and third generations:

$$D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_D & \sin \theta_D \\ 0 & -\sin \theta_D & \cos \theta_D \end{pmatrix}$$
$$L = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_L & \sin \theta_L \\ 0 & -\sin \theta_L & \cos \theta_L \end{pmatrix}.$$

SM-like vector bosons

This model contains vector bosons (VBs) that transform as $(\mathbf{1}, \mathbf{3}, 0)$ under $SU(3)_C \times SU(2)_L \times U(1)_Y$, as in the SM. The coupling is to only third generation. In the gauge basis, the Lagrangian describing the couplings of the VBs to left-handed third-generation fermions is

$$\mathcal{L}_V = g_{qV}^{33} \left(\bar{Q}'_{L3} \gamma^\mu \sigma^I Q'_{L3} \right) V'_\mu + g_{\ell V}^{33} \left(\bar{L}'_{L3} \gamma^\mu \sigma^I L'_{L3} \right) V'_\mu .$$

$$\mathcal{L}_V^{\text{eff}} = -\frac{g_{qV}^{33} g_{\ell V}^{33}}{m_V^2} \left(\bar{Q}'_{L3} \gamma^\mu \sigma^I Q'_{L3} \right) \left(\bar{L}'_{L3} \gamma_\mu \sigma^I L'_{L3} \right) .$$


$$g_1 = 0 \quad , \quad g_2 = -g_{qV}^{33} g_{\ell V}^{33} .$$

The VB model also generates 4 quark and 4 lepton operators that contribute to B_s mixing, $\tau \rightarrow \mu\mu\mu$ e.t.c. Variation of this model with more parameters.


Models: allowed parameter space:

$$R_K \sim \sin \theta_D \cos \theta_D \sin^2 \theta_L$$

VB model: $g_{qV}^{33} = g_{lV}^{33} = \sqrt{0.5}$



U_1 model: $|h_{U_1}^{33}|^2 = 1$



$\tau \rightarrow 3\mu$ (Z' Model)

This decay is particularly interesting because only the VB model contributes to it. The present experimental bound is

$\mathcal{B}(\tau^- \rightarrow \mu^- \mu^+ \mu^-) < 2.1 \times 10^{-8}$ at 90% C.L. . Belle II expects to reduce this limit to $< 10^{-10}$. The reach of LHCb is somewhat weaker, $< 10^{-9}$.

Now, the amplitude for $\tau \rightarrow 3\mu$ depends only on θ_L . The allowed value of θ_L corresponds to the present experimental bound. That is, VB predicts

$$\mathcal{B}(\tau^- \rightarrow \mu^- \mu^+ \mu^-) \simeq 2.1 \times 10^{-8} .$$

Thus, the VB model predicts that $\tau \rightarrow 3\mu$ should be observed at both LHCb and Belle II. This is a smoking-gun signal for the model.

Υ Modes(Leptoquarks)

- $\Upsilon(3S) \rightarrow \mu\tau$:

$$VB \quad \mathcal{B}(\Upsilon(3S) \rightarrow \mu\tau) \simeq 3.0 \times 10^{-9} ,$$

$$U_1 \quad : \quad \mathcal{B}(\Upsilon(3S) \rightarrow \mu\tau)|_{\max} = 8.0 \times 10^{-7} .$$

Belle II should be able to measure $\mathcal{B}(\Upsilon(3S) \rightarrow \mu\tau)$ down to $\sim 10^{-7}$.

- Even though we do not find observable effects in $b \rightarrow s\tau\tau$ or $b \rightarrow s\tau\mu$ others have found larger effects(See for e.g. 1703.09226).

Collider Search: 1706.07808

High- p_T searches are concerned, particularly stringent bounds are set by
 $pp \rightarrow \tau\bar{\tau} + X$

$$\Delta\mathcal{L}_{bb\tau\tau} = -\frac{1}{\Lambda_0^2} (\bar{b}_L\gamma_\mu b_L) (\bar{\tau}_L\gamma_\mu\tau_L) , \quad \Lambda_0^2 = \frac{v^2}{G_1 + G_2} . \quad (10)$$


The present bounds on the EFT scale Λ_0 were derived recasting different ATLAS searches for $\tau\bar{\tau}$ resonances, and read $\Lambda_0 > 0.62$ TeV. Newer fits: $\Lambda_0 \approx 1.2$ TeV, which is well within the experimental limit.

Lepton flavor violating decays: $gg \rightarrow \tau\mu$ (1802.06082, 1802.09822) or
 $gg \rightarrow \bar{t}t\tau\mu$ (1412.7164).

$$\Delta\mathcal{L}_{tt\tau\mu} = -\frac{1}{\Lambda_0^2} (\bar{t}_L\gamma_\mu t_L) (\bar{\tau}_L\gamma_\mu\mu_L) \quad (11)$$

Collider Search: 1706.07808

Z' (1, 3, 0) is strongly constrained (ruled out) unless width is large. Z' (1, 1, 0) explaining only R_K is fine: $M_{Z'} \sim 30$ TeV.



Conclusions

- Several anomalies in B decays indicating lepton non-universal interactions.
- These anomalies may arise from the same New Physics.
- Anomalies indicate LUV. In general we should also observe LFV processes.
- Interesting modes are $\tau \rightarrow 3\mu$ and $\Upsilon(3S) \rightarrow \mu\tau$. Observation of these modes can point to specific models of new physics.
- Other analysis find $b \rightarrow s\tau\tau$ ($B_s \rightarrow \tau^+\tau^-, B \rightarrow M\tau^+\tau^-$) or $b \rightarrow s\tau\mu$ ($B \rightarrow M\tau\mu, B_s \rightarrow \tau\mu$) also promising.