

# Precision measurement with Diboson at the LHC

Da Liu

Argonne National Laboratory


Work in collaboration with **Lian-Tao Wang**

arXiv: 1804.08688

# Motivation


- So far, LHC ( $80 \text{ fb}^{-1}$ ) has not observed any significant excess (Maybe RK?)
- We will have 30 times more data to come.
- Precision measurement enter to a new era at the LHC.
- Some physics scenarios are more sensitive to precision measurement than direct resonance searches.

# Resonance searches reach the slow phase!



Mass scale reach at the LHC

# New physics too heavy to be produced?



# What about LEP?

Compared with LEP, we have more energy and

**Energy helps precision!**

To reach the mass scale  $\Lambda \sim 2 \text{ TeV}$

$$\boxed{\text{LEP}} : \quad \frac{\delta\sigma}{\sigma_{SM}} \sim \frac{m_Z^2}{\Lambda^2} \sim 2.1 \times 10^{-3}$$

$$\boxed{\text{LHC}} : \quad \frac{\delta\sigma}{\sigma_{SM}} \sim \frac{E_c^2}{\Lambda^2} \sim 0.25, \quad E_c \sim 1 \text{ TeV}$$

$$\frac{\delta\sigma}{\sigma_{SM}} \sim \frac{E_c^4}{\Lambda^4} \sim 0.06$$

**LHC has potential.**

**Both interference and energy growing behavior crucial**

# Effective Operators

The model-independent way to capture the new physics effects below the cut-off:

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_{i \in i_6} \frac{c_i}{\Lambda^2} \mathcal{O}_i + \sum_{i \in i_8} \frac{c_i}{\Lambda^4} \mathcal{O}_i$$

- In our parametrization,  $\Lambda$  is the physical cut-off, i.e. the mass of the resonances.
- $c_i$  is the dimensionless Wilson coefficients, which can be large or small (tree or loop) depending the assumptions about the physics at the cut-off.
- Dimension-eight operators maybe relevant in some scenario with:

$$c_8 \gg c_6$$

# Effective Operators

We are focusing on the following dimension-six operators:

$$\begin{aligned}
 \mathcal{O}_W &= \frac{ig}{2} \left( H^\dagger \sigma^a \overleftrightarrow{D}^\mu H \right) D^\nu W_{\mu\nu}^a, & \mathcal{O}_B &= \frac{ig'}{2} \left( H^\dagger \overleftrightarrow{D}^\mu H \right) \partial^\nu B_{\mu\nu} \\
 \mathcal{O}_{2W} &= -\frac{1}{2} D^\mu W_{\mu\nu}^a D_\rho W^{a\rho\nu}, & \mathcal{O}_{2B} &= -\frac{1}{2} \partial^\mu B_{\mu\nu} \partial_\rho B^{\rho\nu} \\
 \mathcal{O}_{HW} &= ig (D^\mu H)^\dagger \sigma^a (D^\nu H) W_{\mu\nu}^a, & \mathcal{O}_{HB} &= ig' (D^\mu H)^\dagger (D^\nu H) B_{\mu\nu} \\
 \mathcal{O}_{3W} &= \frac{1}{3!} g \epsilon_{abc} W_\mu^{a\nu} W_{\nu\rho}^b W^{c\rho\mu}, & \mathcal{O}_T &= \frac{g^2}{2} (H^\dagger \overleftrightarrow{D}^\mu H) (H^\dagger \overleftrightarrow{D}_\mu H) H \\
 \mathcal{O}_R^u &= ig'^2 \left( H^\dagger \overleftrightarrow{D}_\mu H \right) \bar{u}_R \gamma^\mu u_R, & \mathcal{O}_R^d &= ig'^2 \left( H^\dagger \overleftrightarrow{D}_\mu H \right) \bar{d}_R \gamma^\mu d_R \\
 \mathcal{O}_L^q &= ig'^2 \left( H^\dagger \overleftrightarrow{D}_\mu H \right) \bar{Q}_L \gamma^\mu Q_L, & \mathcal{O}_L^{(3)q} &= ig^2 \left( H^\dagger \sigma^a \overleftrightarrow{D}_\mu H \right) \bar{Q}_L \sigma^a \gamma^\mu Q_L
 \end{aligned}$$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \sum_i \frac{c_i}{\Lambda^2} \mathcal{O}_i + \dots$$

# Power Counting of Wilson Coefficients

Model	$\mathcal{O}_{2W}$	$\mathcal{O}_{2B}$	$\mathcal{O}_{3W}$	$\mathcal{O}_{HW}$	$\mathcal{O}_{HB}$	$\mathcal{O}_{W,B}$	$\mathcal{O}_{BB}$
SILH	$\frac{g^2}{g_*^2}$	$\frac{g'^2}{g_*^2}$	$\frac{g^2}{16\pi^2}$	$\frac{g_*^2}{16\pi^2}$	$\frac{g_*^2}{16\pi^2}$	1	$\frac{g^2}{16\pi^2}$
Remedios	1	1	$\frac{g_*}{g}$				
Remedios+MCHM	1	1	$\frac{g_*}{g}$	1	1	1	1
Remedios+ISO(4)	1	1	$\frac{g_*}{g}$	$\frac{g_*}{g}$	1	1	1

- Strongly Interacting Light Higgs (SILH),  $\mathcal{O}_W$  is most relevant one for Di-boson.
- Remedios scenario:  $\mathcal{O}_{2W}$ ,  $\mathcal{O}_{3W}$  enhanced!
- Remedios + ISO(4):  $\mathcal{O}_{HW}$  also enhanced!



# Diboson helicity amplitudes

The helicity amplitudes can be decomposed as:

$$\mathcal{M}_{f_1 \bar{f}_2}^{\lambda_1 \lambda_2, \lambda_3 \lambda_4} = \tilde{\mathcal{M}}_{f_1 \bar{f}_2}^{\lambda_1 \lambda_2, \lambda_3 \lambda_4}(\theta) (\lambda_1 - \lambda_2) (-1)^{\lambda_4} d_{\Delta\lambda_{12}, \Delta\lambda_{34}}^{J_0}$$

with

$$d_{\Delta\lambda_{12}, \Delta\lambda_{34}}^{J_0}, \quad \Delta\lambda_{12} = \lambda_1 - \lambda_2, \quad \Delta\lambda_{34} = \lambda_3 - \lambda_4, \quad J_0 = \max(|\Delta\lambda_{12}|, |\Delta\lambda_{34}|)$$

If  $\tilde{\mathcal{M}}$  is only a function of (s, t, u),  
the amplitude is factorized!

## Explicit formulae for d-functions

$$\begin{aligned} d_{1,2}^2 &= -d_{-1,-2}^2 = \frac{1}{2} \sin \theta (1 + \cos \theta), & d_{1,-2}^2 &= -d_{-1,2}^2 = -\frac{1}{2} \sin \theta (1 - \cos \theta) \\ d_{1,1}^1 &= d_{-1,-1}^1 = \frac{1}{2} (1 + \cos \theta), & d_{1,-1}^1 &= d_{-1,1}^1 = \frac{1}{2} (1 - \cos \theta), \\ d_{1,0}^1 &= -d_{-1,0}^1 = -\frac{\sin \theta}{\sqrt{2}} \end{aligned}$$

# Diboson helicity amplitudes: TT

Subprocess	SM	NP
$u_L \bar{u}_L \rightarrow W_{\pm}^+ W_{\mp}^-$	$-g^2 \frac{s}{2t}$	$-\frac{g^2}{2} \frac{s^2}{\Lambda^4} C_{TWW}$
$d_L \bar{d}_L \rightarrow W_{\pm}^+ W_{\mp}^-$	$-g^2 \frac{s}{2u}$	$-\frac{g^2}{2} \frac{s^2}{\Lambda^4} C_{TWW}$
$u_L \bar{d}_L \rightarrow W_{\pm}^+ Z_{\mp}$	$-\frac{g}{\sqrt{2}} \left( g_Z^{d_L} \frac{s}{t} + g_Z^{u_L} \frac{s}{u} \right)$	$\frac{gg' s_w}{2\sqrt{2}} \frac{s^2}{\Lambda^4} C_{TWB}$
$d_L \bar{u}_L \rightarrow W_{\pm}^- Z_{\mp}$	$-\frac{g}{\sqrt{2}} \left( g_Z^{u_L} \frac{s}{t} + g_Z^{d_L} \frac{s}{u} \right)$	$\frac{gg' s_w}{2\sqrt{2}} \frac{s^2}{\Lambda^4} C_{TWB}$
$u_L \bar{d}_L \rightarrow W_{\pm}^+ \gamma_{\mp}$	$-\frac{g}{\sqrt{2}} \left( g_{\gamma}^{d_L} \frac{s}{t} + g_{\gamma}^{u_L} \frac{s}{u} \right)$	$-\frac{gg' c_w}{2\sqrt{2}} \frac{s^2}{\Lambda^4} C_{TWB}$
$d_L \bar{u}_L \rightarrow W_{\pm}^- \gamma_{\mp}$	$-\frac{g}{\sqrt{2}} \left( g_{\gamma}^{u_L} \frac{s}{t} + g_{\gamma}^{d_L} \frac{s}{u} \right)$	$-\frac{gg' c_w}{2\sqrt{2}} \frac{s^2}{\Lambda^4} C_{TWB}$
$f \bar{f} \rightarrow V_{\pm} V'_{\mp}$	$-g_V^f g_{V'}^f \left( \frac{s}{t} + \frac{s}{u} \right)$	$-\frac{g^2}{4} \frac{s^2}{\Lambda^4} C_{fVV'}^{(8)}$

$\tilde{\mathcal{M}}$

- The transverse WW has t-channel singularity, dominate over LL mode.
  - No interference with dim-6 as angular momentum conservation or helicity selection rules.
- WZ is antisymmetric under (t,u) exchange in the limit of vanishing hypercharge coupling, resulting in the famous amplitude-zero.
- All can interfere with dim-8 operators.

# Diboson helicity amplitudes: LL

Subprocess	SM	NP	$\tilde{\mathcal{M}}$
$u_L \bar{u}_L \rightarrow W_L^+ W_L^-$ $d_L \bar{d}_L \rightarrow Z_L h$	$\frac{1}{\sqrt{2}} \left( \frac{g^2}{2} + \frac{g'^2}{6} \right)$	$\frac{1}{\sqrt{2}} \frac{s}{\Lambda^2} \left( \frac{g^2}{2} c_{qL}^{(3)} + \frac{g'^2}{6} c_{uL}^{(1)} \right)$	
$d_L \bar{d}_L \rightarrow W_L^+ W_L^-$ $u_L \bar{u}_L \rightarrow Z_L h$	$\frac{1}{\sqrt{2}} \left( -\frac{g^2}{2} + \frac{g'^2}{6} \right)$	$\frac{1}{\sqrt{2}} \frac{s}{\Lambda^2} \left( -\frac{g^2}{2} c_{qL}^{(3)} + \frac{g'^2}{6} c_{dL}^{(1)} \right)$	
$u_R \bar{u}_R \rightarrow W_L^+ W_L^-$	$\frac{\sqrt{2}}{3} g'^2$	$\frac{\sqrt{2}}{3} g'^2 \frac{s}{\Lambda^2} c_{uR}^{(1)}$	
$u_R \bar{u}_R \rightarrow W_L^+ W_L^-$	$\frac{\sqrt{2}}{3} g'^2$	$\frac{\sqrt{2}}{3} g'^2 \frac{s}{\Lambda^2} c_{uR}^{(1)}$	
$d_R \bar{d}_R \rightarrow W_L^+ W_L^-$	$-\frac{\sqrt{2}}{6} g'^2$	$-\frac{\sqrt{2}}{6} g'^2 \frac{s}{\Lambda^2} c_{dR}^{(1)}$	
$u_L \bar{d}_L \rightarrow W_L^+ Z_L(h)$ $d_L \bar{u}_L \rightarrow W_L^- Z_L(h)$	$-\frac{g^2}{2}$	$-\frac{g^2}{2} \frac{s}{\Lambda^2} c_{qL}^{(3)}$	

$$c_{qL}^{(3)} = c_W + c_{HW} - c_{2W} + 4c_L^{(3)q}$$

$$c_{uL}^{(1)} = c_B + c_{HB} - c_{2B} + 4c_L^q$$

$$c_{dL}^{(1)} = c_B + c_{HB} - c_{2B} - 4c_L^q$$

$$c_{uR}^{(1)} = c_B + c_{HB} - c_{2B} + 3c_{uR}$$



$$c_{dR}^{(1)} = c_B + c_{HB} - c_{2B} - 6c_{dR}$$

- Because of the smallness of  $U(1)_Y$  coupling, the dominant contribution comes from the combination  $c_{qL}^{(3)}$
- Wh Zh no transverse backgrounds, but suffer from W+jets and Z+jets.

$$\mathcal{O}_{3W} = \frac{1}{3!} g \epsilon_{abc} W_{\mu}^{a\nu} W_{\nu\rho}^b W^{c\rho\mu},$$

- Helicity selection rules tells us this only contribute to ++ helicity final states (see Azatov et al.)
 
$$|h(A_3)| = 1 - [g].$$
- Underlying SUSY ward identity for gauge interactions at tree level ensure ++ SM amplitudes are zero (the same reason as the vanishing of the ++++ four gluon amplitudes)
- Interference with SM not growing with energy, hard to probe at the hadron collider
- Can be improved by explore the azimuthal angles of decay products (See Azatov et al. and Panico et al.)

# WW and WZ at the LHC



**Focusing on the central region improves the significance**

# We are focusing on the semi-leptonically decaying channels

$$\begin{aligned} pp \rightarrow WV \rightarrow \ell\nu q\bar{q}, & \quad \text{BR}(W^+W^- \rightarrow \ell\nu q\bar{q}) = 29.2\%, & \quad \text{BR}(W^\pm Z \rightarrow \ell\nu q\bar{q}) = 15.1\% \\ pp \rightarrow Wh \rightarrow \ell\nu b\bar{b}, & \quad \text{BR} = 12.6\% \\ pp \rightarrow Zh \rightarrow \ell^+\ell^- b\bar{b}, & \quad \text{BR} = 3.92\% \\ pp \rightarrow Zh \rightarrow \nu\bar{\nu} b\bar{b}, & \quad \text{BR} = 11.6\% \end{aligned}$$



- Compared with fully leptonic channels, these have one order of magnitude larger rate.
- Although suffered from reducible backgrounds  $V + \text{jets}$ , a lot of data can make big difference here!
- Jet substructure methods play an important role to suppress the reducible backgrounds.
- 13 TeV  $W$ -jet tagging has been improved by a factor of 2 with 8 TeV.

# Polarization tagging

$$P_+ = \frac{3}{8}(1 - \cos \theta^*)^2, \quad P_- = \frac{3}{8}(1 + \cos \theta^*)^2, \quad P_0 = \frac{3}{4} \sin^2 \theta^*$$

$\cos \theta^*$  can be determined by the momenta of the leptons and neutrinos


$$\cos \theta^* = \frac{E_\ell - E_\nu}{|\vec{p}_\ell + \vec{p}_\nu|}$$



$$\epsilon_L \equiv \epsilon_{p_{T,\eta}}^L \times \epsilon_{\cos \theta^*}^L = 0.75,$$

$$\epsilon_T \equiv \epsilon_{p_{T,\eta}}^T \times \epsilon_{\cos \theta^*}^T = 0.5.$$


# Reach in different energy bins



With HL, weakly coupled EFT consistent!




# Bounds on O3W



- Bounds are in tension with weakly coupled EFT.
- Still useful for strongly coupled case, e.g. Strong multi-pole interactions.

Bound at LHC14,  $\Delta_{\text{sys}} \in [3\%, 10\%]$ ,  $c_i = 1$



**Bounds from other measurements**

- $O_W + O_B$ , LEP S-parameter
- $O_{HW} - O_{HB}$ , HL-LHC  $h \rightarrow Z \gamma$
- $O_L^{(3)q}$  LEP  $\delta g_{Zb_L b_L}$

# Mass scale reach in different scenarios

Model	Di-boson	S-parameter	LHC $h \rightarrow Z\gamma$	LHC $h \rightarrow \gamma\gamma$	LHC dilepton
SILH	4.0	2.5	$1.7\sqrt{\frac{g_*}{4\pi}}$	0.34	$0.69\sqrt{\frac{4\pi}{g_*}}$
Remedios	$10.6\sqrt{\frac{g_*}{4\pi}}$				13.4
Remedios+MCHM	$10.6\sqrt{\frac{g_*}{4\pi}}$	2.5	1.7	6.5	13.4
Remedios+ISO(4)	$17.6\sqrt{\frac{g_*}{4\pi}}$	2.5	$7.5\sqrt{\frac{g_*}{4\pi}}$	6.5	13.4

- Strongly Interacting Light Higgs (SILH), Di-boson can beat the LEP EWPT.
- Pure Strong multi-pole interactions, LHC dilepton bounds on O2W maybe the most relevant ones.
- Remedios+ISO(4), Di-boson can be better for large coupling  $g^* > 7$ .

# Conclusion

- New physics may only show its tail at the LHC, it is important to do the precision measurement.
- The EFT is a convenient and model-independent way to capture the effects.
- With high energy at hand, LHC can beat LEP precision.
- Non-resonant, broad features. Difficult. But a lot data can make a significant difference here!

Back-up Slides

# Helicity structure for WW

$$q_L \bar{q}_R \rightarrow W^+ W^-$$

$(h_{W^+}, h_{W^-})$	SM	$\mathcal{O}_W$	$\mathcal{O}_{HW}$	$\mathcal{O}_B$	$\mathcal{O}_{HB}$	$\mathcal{O}_{3W}$
$(\pm, \mp)$	1	0	0	0	0	0
$(0, 0)$	1	$\frac{E^2}{\Lambda^2}$	$\frac{E^2}{\Lambda^2}$	$\frac{E^2}{\Lambda^2}$	$\frac{E^2}{\Lambda^2}$	0
$(0, \pm), (\pm, 0)$	$\frac{m_W}{E}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$
$(\pm, \pm)$	$\frac{m_W^2}{E^2}$	$\frac{m_W^2}{\Lambda^2}$	$\frac{m_W^2}{\Lambda^2}$	$\frac{m_W^2}{\Lambda^2}$	0	$\frac{E^2}{\Lambda^2}$

$$q_R \bar{q}_L \rightarrow W^+ W^-$$

$(h_{W^+}, h_{W^-})$	SM	$\mathcal{O}_W$	$\mathcal{O}_{HW}$	$\mathcal{O}_B$	$\mathcal{O}_{HB}$	$\mathcal{O}_{3W}$
$(\pm, \mp)$	0	0	0	0	0	0
$(0, 0)$	1	$\frac{m_W^2}{\Lambda^2}$	$\frac{m_W^2}{\Lambda^2}$	$\frac{E^2}{\Lambda^2}$	$\frac{E^2}{\Lambda^2}$	0
$(0, \pm), (\pm, 0)$	$\frac{m_W}{E}$	$\frac{m_W^2 m_Z^2}{\Lambda^2 E^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{Em_W}{\Lambda^2}$	$\frac{m_W^2 m_Z^2}{\Lambda^2 E^2}$
$(\pm, \pm)$	$\frac{m_W^2}{E^2}$	$\frac{m_W^2}{\Lambda^2}$	$\frac{m_W^2}{\Lambda^2}$	$\frac{m_W^2}{\Lambda^2}$	0	$\frac{m_W^2}{\Lambda^2}$

- Whether interference or not depends on polarization of WW. Polarization differentiation can be crucial.

# Operator relations

- Equation of motion or field redefinition

$$D^\nu W_{\mu\nu}^a = igH^\dagger \frac{\sigma^a}{2} \overleftrightarrow{D}_\mu H + g \sum_f \bar{f}_L \frac{\sigma^a}{2} \gamma_\mu f_L,$$

$$\partial^\nu B_{\mu\nu} = ig' Y_H H^\dagger \overleftrightarrow{D}_\mu H + g' \sum_f \left[ Y_L^f \bar{f}_L \gamma_\mu f_L + Y_R^f \bar{f}_R \gamma_\mu f_R \right]$$

$$D^\nu G_{\mu\nu}^A = g_s \sum_q \bar{q} T^A \gamma_\mu q,$$

- Partial integral

$$\begin{aligned} \mathcal{O}_B &= \mathcal{O}_{HB} + \frac{1}{4} \mathcal{O}_{BB} + \frac{1}{4} \mathcal{O}_{WB}, & \mathcal{O}_{WW} &= g^2 |H|^2 W_{\mu\nu}^a W^{\mu\nu a} \\ \mathcal{O}_W &= \mathcal{O}_{HW} + \frac{1}{4} \mathcal{O}_{WW} + \frac{1}{4} \mathcal{O}_{WB}. & \mathcal{O}_{WB} &= g' g H^\dagger \sigma^a H W_{\mu\nu}^a B^{\mu\nu} \end{aligned}$$

- We can use them to rewrite the more derivative operators with more fields  $\rightarrow$  Warsaw basis.

$$\mathcal{O}_W = -\frac{3}{2} \mathcal{O}_H + 2\mathcal{O}_6 + \frac{1}{2} (\mathcal{O}_{y_u} + \mathcal{O}_{y_d} + \mathcal{O}_{y_e}) + \frac{1}{4} \sum_f \mathcal{O}_L^{(3)f}$$

$$\mathcal{O}_B = -\frac{1}{2} \mathcal{O}_T + \frac{1}{2} \sum_f \left( Y_L^f \mathcal{O}_L^f + Y_R^f \mathcal{O}_R^f \right)$$

see for example Elias-Miro et al.

# Phenomenology of the effective operators

## hVV contact interactions, $V = W, Z, A$

$$\mathcal{L}_h = \frac{2m_W^2}{\Lambda^2} \left( \frac{h}{v} + \frac{h^2}{2v^2} \right) \left\{ (\hat{c}_W W_\mu^- \mathcal{D}^{\mu\nu} W_\nu^+ + h.c.) + Z_\mu \mathcal{D}^{\mu\nu} \left[ \hat{c}_Z Z_\nu + \left( \frac{2\hat{c}_W}{\sin 2\theta_W} - \frac{\hat{c}_Z}{\tan \theta_W} \right) A_\nu \right] \right\} \\ - 4 \frac{m_W^2}{\Lambda^2} \left( \frac{h}{v} + \frac{h^2}{2v^2} \right) \left\{ \frac{c_{HW}}{2} W^{+\mu\nu} W_{\mu\nu}^- + \frac{c_{HW} + \tan^2 \theta_W c_{HB}}{4} Z^{\mu\nu} Z_{\mu\nu} - 2 \sin^2 \theta_W c_{\gamma Z} A^{\mu\nu} Z_{\mu\nu} \right\}$$

$$\mathcal{D}^{\mu\nu} = \partial^\mu \partial^\nu - \square \eta^{\mu\nu}, \quad \hat{c}_W = c_W + c_{HW}, \quad \hat{c}_Z = \hat{c}_W + \tan^2 \theta_W (c_B + c_{HB}), \quad c_{\gamma Z} = -\frac{c_{HW} - c_{HB}}{4 \sin 2\theta_W}$$

- OW and OB don't contribute to on-shell photon production
- OHW and OHB contribute to  $hZ\gamma$ , loop level generated in minimal coupled theory



# Phenomenology of the effective operators

TGC:

$$\mathcal{L}_V = -\frac{\tan \theta_W}{2} \hat{S} W_{\mu\nu}^{(3)} B^{\mu\nu} + ig \cos \theta_W \delta g_1^Z (W_{\mu\nu}^+ W^{-\mu} - W_{\mu\nu}^- W^{+\mu}) Z^\nu$$

$$+ ig (\cos \theta_W \delta \kappa_Z Z^{\mu\nu} + \sin \theta_W \delta \kappa_\gamma A^{\mu\nu}) W_\mu^+ W_\nu^- + ig \cos \theta_W \frac{\lambda_Z}{m_W^2} W_{\lambda\mu}^+ W^{-\mu}_\nu Z^{\nu\lambda} + ie \frac{\lambda_\gamma}{m_W^2} W_{\lambda\mu}^+ W^{-\mu}_\nu A^{\nu\lambda}$$

$$\hat{S} = (c_W + c_B) \frac{m_W^2}{\Lambda^2}, \quad \delta g_1^Z = -\frac{c_W + c_{HW}}{\cos^2 \theta_W} \frac{m_W^2}{\Lambda^2}, \quad \delta \kappa_\gamma = -(c_{HW} + c_{HB}) \frac{m_W^2}{\Lambda^2}$$

$$\delta \kappa_Z = \delta g_1^Z - \tan^2 \theta_W \delta \kappa_\gamma, \quad \lambda_Z = \lambda_\gamma = c_{3W} \frac{m_W^2}{\Lambda^2}$$

- OW and OB contribute to the S parameter, constrained by LEP
- O3W modify the magnetic dipole and to the electric quadrupole of the W, can only arise from loop level in minimal coupled theory

# Phenomenology of the effective operators

## Fermion couplings:

$$\mathcal{L}_f = \frac{4m_W^2}{\Lambda^2} \left(1 + \frac{h}{v}\right)^2 \left[ \frac{g}{\sqrt{2}} c_L^{(3)q} W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c. \right. \\ \left. - \frac{g}{2c_w} Z_\mu \left( c_R^u \bar{u}_R \gamma^\mu u_R + c_R^d \bar{d}_R \gamma^\mu d_R + (c_L^q - c_L^{(3)q}) \bar{u}_L \gamma^\mu u_L + (c_L^q + c_L^{(3)q}) \bar{d}_L \gamma^\mu d_L \right) \right]$$

- Contribute to the Z-pole variables.
- We will consider the flavour-universal effects
- Contact interactions between ffVh, longitudinal modes growing with energy.

# Phenomenology of the effective operators

## Dimension-eight:

$$\begin{aligned}
 \mathcal{L}_8 = & \frac{1}{\Lambda^4} T_f^{\mu\nu} \left[ c_{TWW} g^2 (W_{\mu\rho}^+ W_{\nu}^{-\rho} + W_{\nu\rho}^+ W_{\mu}^{-\rho}) + (c_{TWW} g^2 c_w^2 + c_{TBB} g'^2 s_w^2 - 2T_f^3 c_{TWB} g g' s_w c_w) Z_{\mu\rho} Z_{\nu}^{\rho} \right. \\
 & + (c_{TWW} + c_{TBB} + 2T_f^3 c_{TWB}) e^2 A_{\mu\rho} A_{\nu}^{\rho} + (2c_w s_w (c_{TWW} g^2 - c_{TBB} g'^2) + 2T_f^3 c_{TWB} g g' (c_w^2 - s_w^2)) Z_{\mu\rho} A_{\nu}^{\rho} \\
 & \left. + \frac{1}{2} (c_{TH} - 2T_f^3 c_{TH}^{(3)}) g^2 (\partial_{\mu} h \partial_{\nu} h + m_Z^2 Z_{\mu} Z_{\nu}) + (c_{TH} + 2T_f^3 c_{TH}^{(3)}) g^2 m_W^2 W_{\mu}^{-} W_{\nu}^{+} \right] \\
 & + c_{TWB} \frac{\sqrt{2} g g'}{\Lambda^4} \left( T_{fL}^{+\mu\nu} W_{\mu\rho}^{+} + T_{fL}^{-\mu\nu} W_{\mu\rho}^{-} \right) (-s_w Z_{\nu}^{\rho} + c_w A_{\nu}^{\rho}) \\
 & + c_{TH}^{(3)} \frac{\sqrt{2} g^2 m_W}{\Lambda^4} \left( T_{fL}^{+\mu\nu} W_{\mu}^{+} (-i\partial_{\nu} h - m_Z Z_{\nu}) + T_{fL}^{-\mu\nu} W_{\mu}^{-} (i\partial_{\nu} h - m_Z Z_{\nu}) \right)
 \end{aligned}$$

– New signatures:  $ZZ, \gamma\gamma$

– Interference with SM:  $\frac{\delta\sigma}{\sigma_{\text{SM}}} \sim \frac{E^4}{\Lambda^4}$

– Can be enhanced by strong coupling