

Contribution ID: 140

Type: Oral

In-beam and decay spectroscopy of ²⁵¹Md

Monday, 13 June 2022 11:50 (20 minutes)

Experiments on the heaviest nuclei are addressing the fundamental issue of the maximum limits of nuclear mass and charge. The difficulty of producing such exotic nuclei, however, creates an inherent challenge for elucidating the detailed structure of these extreme nuclear systems. By contrast, nuclei in the vicinity of Z = 100 and N = 152 may be studied in much greater detail, and yet they involve intrinsic excitations of the same single-particle orbits that lie close to the Fermi surface of the spherical superheavy nuclei-[1]. In particular, the structure of the low-lying energy levels in odd-mass transfermium nuclei is dominated by the unpaired particle, which gives direct information about the orbitals involved. The odd-Z nucleus ²⁵¹Md has recently been the subject of considerable interest in this regard, with the observation of states based on the $\frac{1}{2}$ [521] Nilsson level-[2], the $\frac{7}{2}$ [514] Nilsson level-[3], as well as a high-K isomer-[4]. These experiments are expanding our view of not only the transfermium region, but the superheavy nuclei as well.

We have performed a new experiment to study 251 Md via both in-beam and decay spectroscopy. The 205 Tl(48 Ca,2n) fusion-evaporation reaction was used to populate excited states in 251 Md. The newly commissioned Argonne Gas-Filled Analyzer (AGFA) was used to separate the unreacted beam and the 251 Md recoils, which were implanted into a 160×160 pixel silicon double-sided strip detector (DSSD) at the focal plane of AGFA. Prompt γ rays were detected with Gammasphere and correlated with 251 Md based on their time-of-flight and implantation energy in the AGFA focal plane. Long-lived isomers were detected by the observation of bursts of conversion electrons in the same DSSD pixel as an implantation event, and delayed γ rays emitted after isomer-decay events were detected with the X-array, which was arranged around the DSSD. With this setup, we have observed many new transitions in both the prompt and the delayed γ spectra, including transitions above the isomeric state. These results and their implications for the structure of 251 Md will be presented.

- [1] R.-D.~Herzberg \textit{et al.}, Nature 442, 896 (2006)\\
- [2] A.~Chatillon \textit{et al.}, Phys. Rev. Lett. 98, 132503 (2007)\\
- [3] R.~Briselet \textit{et al.}, Phys. Rev. C 102, 014307 (2020)\\
- [4] T.~Goigoux \textit{et al.}, Eur. Phys. J. A 57, 321 (2021)\\

Primary authors: MORSE, C. (National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973); HUANG, T. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); SEWERYNIAK, D. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); CLARK, R.M. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); BACK, B.B. (Physics Division, Argonne National Laboratory, Berkeley, CA 94720); BACK, B.B. (Physics Division, Argonne National Laboratory, Berkeley, CA 94720); CAMPBELL, C.M. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); CARPENTER, M.P. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); CHOWDHURY, P. (Department of Physics and Applied Physics, University of Massachusetts Lowell, Lowell, MA 01854); CRAWFORD, H.L. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); FALLON, P. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); FALLON, P. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); FAVIER, Z. (Irfu, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France); KHOO, T.L. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); KONICHI, A. (IJCLab, IN2P3-CNRS Orsay, France); LAURITSEN, T. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); MACCHIAVELLI, A.O. (Nuclear Science Division, Argonne National Laboratory, Argonne, IL 60439); MACCHIAVELLI, A.O. (Nuclear Science Division, Argonne, IL 60439); MACCHIAVELLI, A.O. (Nuclear Science Division, Magonne, Macchiavell, A.O. (Nuclear Science Division, Argonne, IL 60439); MACCHIAVELLI, A.O. (

Lawrence Berkeley National Laboratory, Berkeley, CA 94720); POTTERVELD, D. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); REVIOL, W. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); RUDOLPH, D. (Department of Physics, Lund University, 22100 Lund, Sweden); SANTAMARIA, C. (Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720); SAVARD, G. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); STOLZE, S. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); WILSON, G.L. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); WU, J. (Physics Division, Argonne National Laboratory, Argonne, IL 60439); ZHU, S. (Physics Division, Argonne National Laboratory, Argonne, IL 60439)

Presenter: MORSE, C. (National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973)

Session Classification: NS2022 Plenary

Track Classification: Oral Presentations